

Constituent monopoles without gauge fixing

Thomas C. Kraan^a and Pierre van Baal^{a*}

^aInstituut-Lorentz for Theoretical Physics, University of Leiden,
 PO Box 9506, NL-2300 RA Leiden, The Netherlands.

We discuss the recent construction of new exact finite temperature instanton solutions with a non-trivial value of the Polyakov loop at infinity. They can be shown, in a precise and gauge invariant way, to be formed by the superposition of n BPS monopoles for an $SU(n)$ gauge group.

1. Introduction

Instantons at finite temperature (or calorons) are constructed on $\mathbb{R}^3 \times S^1$, taking a periodic array of instantons. For $SU(2)$ the five parameter Harrington-Shepard solution [1] can be formulated within the 't Hooft ansatz. New exact solutions with a non-trivial value of the Polyakov loop at infinity [2] were only constructed very recently, either using [3] results due to Nahm [4] or by using [5] the well-known ADHM construction [6], translated by Fourier transformation to the Nahm language. Thus mapped to an Abelian problem on the circle, the quadratic ADHM constraint is solved [5].

2. New caloron solutions

In the periodic gauge, $A_\mu(x + \beta) = A_\mu(x)$, the Polyakov loop at spatial infinity

$$\mathcal{P}_\infty = \lim_{|\vec{x}| \rightarrow \infty} P \exp\left(\int_0^\beta A_0(\vec{x}, t) dt\right), \quad (1)$$

after a constant gauge transformation, is characterised by $(\sum_{m=1}^n \mu_m = 0)$

$$\mathcal{P}_\infty^0 = \exp[2\pi i \text{diag}(\mu_1, \dots, \mu_n)], \quad (2)$$

$$\mu_1 < \dots < \mu_n < \mu_{n+1} \equiv \mu_1 + 1.$$

A non-trivial value, $\mathcal{P} \notin Z_n$, acts like a Higgs field. We found [5c] a remarkably simple formula for the action density, valid for arbitrary $SU(n)$. Using the classical scale invariance to put $\beta = 1$,

$$\text{tr} F_{\mu\nu}^2 = \partial_\mu^2 \partial_\nu^2 \log \psi, \quad \psi = -\cos(2\pi t) + \frac{1}{2} \text{tr} \prod_{m=1}^n A_m,$$

* Presented by second author.

$$A_m \equiv \frac{1}{r_m} \begin{pmatrix} r_m & |\vec{y}_m - \vec{y}_{m+1}| \\ 0 & r_{m+1} \end{pmatrix} \begin{pmatrix} c_m & s_m \\ s_m & c_m \end{pmatrix}, \quad (3)$$

with $r_m = |\vec{x} - \vec{y}_m|$ the center of mass radius of the m^{th} constituent monopole, which can be assigned a mass $16\pi^2 \nu_m$, where $\nu_m \equiv \mu_{m+1} - \mu_m$. Also $r_{n+1} \equiv r_1$, $\vec{y}_{n+1} \equiv \vec{y}_1$, $c_m \equiv \cosh(2\pi \nu_m r_m)$ and $s_m \equiv \sinh(2\pi \nu_m r_m)$. The order of matrix multiplication is crucial here, $\prod_{m=1}^n A_m \equiv A_n \dots A_1$.




Figure 1. Action densities for the $SU(3)$ caloron on equal logarithmic scales, cut off at $1/e$, for $t = 0$ in the plane defined by $\vec{y}_1 = (-\frac{1}{2}, \frac{1}{2}, 0)$, $\vec{y}_2 = (0, \frac{1}{2}, 0)$ and $\vec{y}_3 = (\frac{1}{2}, -\frac{1}{2}, 0)$, in units of β , for $\beta = 1/4, 1/3$ and $2/3$ from top to bottom, using $(\mu_1, \mu_2, \mu_3) = (-17, -2, 19)/60$.

For $\mathcal{P}_\infty = \exp(2\pi i \omega \tau_3)$ the $SU(2)$ gauge field reads [5a], in terms of the anti-selfdual 't Hooft tensor $\tilde{\eta}$ and Pauli matrices τ_a ,

$$A_\mu(x) = \frac{i}{2} \tilde{\eta}_{\mu\nu}^3 \tau_3 \partial_\nu \log \phi(x) + \frac{i}{2} \phi(x) \operatorname{Re}((\tilde{\eta}_{\mu\nu}^1 - i \tilde{\eta}_{\mu\nu}^2)(\tau_1 + i \tau_2) \partial_\nu \chi(x)), \quad (4)$$

where $\phi^{-1} = 1 - \frac{\pi \rho^2}{\psi} \left(\frac{s_1 c_2}{r_1} + \frac{s_2 c_1}{r_2} + \frac{\pi \rho^2 s_1 s_2}{r_1 r_2} \right)$ and $\chi = \frac{\pi \rho^2}{\psi} \left(e^{-2\pi i t} \frac{s_1}{r_1} + \frac{s_2}{r_2} \right) e^{2\pi i \nu_1 t}$, with $\nu_1 = 2\omega$, $\nu_2 = 1 - 2\omega$ and $\pi \rho^2 = |\tilde{y}_2 - \tilde{y}_1|$. The solution is presented in the “algebraic” gauge, $A_\mu(x + \beta) = \mathcal{P}_\infty A_\mu(x) \mathcal{P}_\infty^{-1}$.




Figure 2. Action densities for the $SU(2)$ caloron on equal logarithmic scales, cut off below $1/e^2$, for $t = 0$, $\omega = 0.125$, $\beta = 1$ and $\rho = 1.6, 1.2$ and 0.8 (from top to bottom).




Figure 3. As in fig. 2, now cut off below $1/e$, for $t = 0$, $\rho = \beta = 1$ with $\omega = \frac{1}{4}$ (top) and 0 (bottom).

For small ρ , equivalent to large β , the caloron approaches the ordinary single instanton solution, with no dependence on \mathcal{P}_∞ . Finite size effects set in roughly when $\rho = \frac{1}{2}\beta$. At this point, for $\nu_i \neq 0$, two lumps (n for $SU(n)$) are formed, whose separation grows as $\pi \rho^2 / \beta$. When $\mathcal{P}_\infty = (-1)$ for $SU(2)$, one of the lumps disappears, as $\nu_{1(2)} = 0$, and the spherically symmetric Harrington-Shepard solution is retrieved.

A non-trivial value of \mathcal{P}_∞ will modify the vacuum fluctuations and thereby leads to a non-zero vacuum energy density [2] as compared to $\mathcal{P} \in \mathbb{Z}_n$. A dilute, semi-classical instanton calculation is no longer considered a reliable starting point for QCD. Rather, it is the monopole constituent nature from which we should draw important lessons for QCD [5b].

3. Monopoles from instantons

At small β the solution becomes static and the lumps are well separated and spherically symmetric. As they are self-dual, they are necessarily BPS monopoles [7]. Also, when sending (at fixed β) one of the constituents to infinity, $|\tilde{y}_m| \rightarrow \infty$, the solution becomes static and yields a simple way to obtain $SU(n)$ monopole solutions [5c]. Explicitly we find (assuming $\nu_n \neq 0$) in the limit $|\tilde{y}_n| \rightarrow \infty$, which removes the n -th constituent,

$$A_n \rightarrow 2c_n \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}, \quad A_{n-1} \rightarrow \frac{|\tilde{y}_n|}{r_{n-1}} \begin{pmatrix} s_{n-1} & c_{n-1} \\ s_{n-1} & c_{n-1} \end{pmatrix}, \quad (5)$$

implying $\psi(x) \rightarrow 2|\tilde{y}_n| \exp(2\pi \nu_n |\tilde{y}_n - \tilde{x}|) \tilde{\psi}(\tilde{x})$, with

$$\tilde{\psi}(\tilde{x}) = \frac{1}{2} \operatorname{tr} \left\{ \frac{1}{r_{n-1}} \begin{pmatrix} s_{n-1} & c_{n-1} \\ 0 & 0 \end{pmatrix} \prod_{m=1}^{n-2} A_m \right\}. \quad (6)$$

As was emphasised in ref. [5c], the energy density of the $SU(n)$ monopole is easily found from eq. (3) (for a detailed description of some special cases see ref. [8])

$$\mathcal{E}(\tilde{x}) = -\frac{1}{2} \operatorname{tr} F_{\mu\nu}^2(\tilde{x}) = -\frac{1}{2} \Delta^2 \log \tilde{\psi}(\tilde{x}). \quad (7)$$

4. Instantons from monopoles

The new caloron solutions provide examples of gauge fields with topological charge built out of monopole fields, a construction going back to

Taubes [9]. Non-trivial $SU(2)$ monopole fields are classified by the winding number of maps from S^2 to $SU(2)/U(1) \sim S^2$, where $U(1)$ is the unbroken gauge group. Isospin orientations for a configuration made out of monopoles with opposite charges behave as shown in fig. 4 (top), in a suitable gauge and sufficiently far from the core of both monopoles. Taubes constructed topologically non-trivial configurations by creating a monopole anti-monopole pair, bringing them far apart, rotating one of them over a full rotation and finally bringing them together to annihilate (cmp. fig. 5). We can describe this as a closed monopole line (or loop) with the orientation of the core defined by $SU(2)/U(1) \sim S^2$, “twisting” along the loop, thus describing a Hopf fibration [5b] (see fig. 4 (bottom)). The only topological invariant available to characterise the homotopy type of this Hopf fibration is the Pontryagin index. It prevents full annihilation of the “twisted” monopole loop.

For large ρ , eq. (4) gives up to exponential corrections, i.e. outside the cores of the constituents,

$$A_\mu = \frac{i}{2} \tau_3 \bar{\eta}_{\mu\nu}^3 \partial_\nu \log \phi_0, \quad \phi_0 \equiv \frac{r_1 + r_2 + \pi \rho^2}{r_1 + r_2 - \pi \rho^2}. \quad (8)$$

This describes two Abelian Dirac monopoles and one easily verifies $\log \phi_0$ is harmonic, as required by self-duality. Furthermore ϕ_0^{-1} vanishes on the line connecting the two monopole centers, giving rise to return flux, absent in the full theory. The relative phase $e^{-2\pi i t}$ in the expression for χ given




Figure 4. Topological charge constructed from oppositely charged monopoles by rotating one of them. For a closed monopole line, the embedding of the unbroken subgroup makes a full rotation.

before, describes the full rotation of the core of a constituent monopole, required so as to give rise to non-trivial topology.

A conjectured QCD application, in the form of a hybrid monopole-instanton liquid, was discussed in ref. [5b]. Abelian projection applied to our solutions was also discussed at this conference [10].




Figure 5. Action density in the z - t -plane for $x=y=0$, $\omega=\frac{1}{4}$, $\rho=\frac{1}{2}$ and $\beta=1$ on a linear scale. One can trace the constituent monopoles in the low field regions, “annihilating” to give an instanton.

REFERENCES

1. B.J. Harrington and H.K. Shepard, Phys. Rev. **D17** (1978) 2122; **D18** (1978) 2990
2. D.J. Gross, R.D. Pisarski and L.G. Yaffe, Rev. Mod. Phys. **53** (1983) 43
3. K. Lee, Phys. Lett. **B426** (1998) 323; K. Lee and C. Lu, Phys. Rev. **D58** (1998) 025011
4. W. Nahm, *Self-dual monopoles and calorons*, in: Lecture Notes in Physics 201 (1984) 189
5. a: T.C. Kraan and P. van Baal, Phys. Lett. **B428** (1998) 268; b: hep-th/9805168, hep-th/9805201; c: hep-th/9806034
6. M.F. Atiyah, N.J. Hitchin, V. Drinfeld and Yu.I. Manin, Phys. Lett. **65A** (1978) 185
7. E.B. Bogomol'ny, Sov. J. Nucl. **24** (1976) 449; M.K. Prasad and C.M. Sommerfield, Phys. Rev. Lett. **35** (1975) 760
8. C. Lu, hep-th/9806237.
9. C. Taubes, in: *Progress in gauge field theory*, eds. G.'t Hooft e.a., (Plenum Press, New York, 1984) p.563
10. Talks by R. Brower and J. Negele at this conference